

FORAGE SUITABILITY GROUP SAND

FSG No.: G063AY300SD

Major Land Resource Area: 63A - Northern Rolling Pierre Shale Plains

Physiographic Features

The soils in this group are found on flood plains, terraces, and upland slopes.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	1300	2950
Slope (percent):	0	6
Flooding:		
Frequency:	None	Frequent
Duration:	None	Brief
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Negligible	Very low

Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 63A. Average annual precipitation for all climate stations listed below is about 17 inches. About 77 percent of that occurs during the months of April through September. On average, there are about 25 days with greater than .1 inches of precipitation during that same time period.

Average annual snowfall ranges from 24 inches at Midland to 48 inches at Milesville. Snow cover at depths greater than 1 inch range from 27 days at Midland to 82 days at Timber Lake.

Average July temperatures across the MLRA are about 75°F and average January temperatures are about 17°F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -37 at Kennebec and a high of 114 at both Kennebec and Midland. The MLRA lies in USDA Plant Hardiness Zones 4a, 4b, and 5a.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data, access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	129	162
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 20	May 04
Last Frost in Spring (32 deg): (1 year in 10 later than)	May 31	May 16
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Sep 09	Sep 24
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 17	Oct 01
Length of Growing Season (32 deg) (days): (9 years in 10 at least)	110	139

	From	To
Growing Degree Days (40 deg):	4442	5149
Growing Degree Days (50 deg):	2517	3083
Annual Minimum Temperature:	-30	-15
Mean annual precipitation (inches):	16	18

Monthly precipitation (inches) and temperature (F)

2 years in 10:	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Less Than	0.10	0.09	0.31	0.82	1.44	1.55	0.90	0.64	0.41	0.30	0.08	0.16
Precip. More Than	0.60	0.79	2.37	3.46	3.82	4.55	3.58	2.46	1.98	2.06	1.07	0.91
Monthly Average:	0.30	0.42	1.20	1.99	2.86	3.06	2.23	1.80	1.31	1.12	0.48	0.45
Temp. Min.	3.9	9.7	20.0	32.9	44.0	53.9	59.6	57.3	46.4	35.5	20.8	7.9
Temp. Max.	32.8	38.6	48.3	63.1	74.1	83.8	92.2	90.6	79.3	66.4	48.4	35.9
Temp. Avg.	17.1	22.9	33.0	46.7	58.0	68.0	75.0	73.0	61.7	49.6	33.5	20.5

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
SD5691	Mobridge SD	1961	1990
SD8307	Timber Lake SD	1961	1990
SD6170	Oahe Dam SD	1961	1990
SD5506	Midland SD	1961	1990
SD5544	Milesville SD	1961	1990
SD6552	Philip SD	1961	1990
SD5891	Murdo SD	1961	1990
SD4516	Kennebec SD	1961	1990

Soil Interpretations

This group consists of very deep, somewhat excessively to excessively drained, coarse textured soils formed in sandy eolian and alluvial materials. Permeability is rapid.

Drainage Class:	Somewhat excessively drained	To	Excessively drained
Permeability Class:	None selected	To	None selected
(0 - 40 inches)			
Frost Action Class:	Low	To	Low

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent):	0.5	2.0
(surface layer)		
Electrical Conductivity (mmhos/cm):	0	4
(0 - 24 inches)		
Sodium Absorption Ratio:	0	0
(0 - 12 inches)		
Soil Reaction (1:1) Water (pH):	5.6	8.4
(0 - 12 inches)		
Available Water Capacity (inches):	3	6
(0 - 60 inches)		
Calcium Carbonate Equivalent	0	18
(0 - 12 inches)		

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed at <http://plants.usda.gov/>.

Sand

Pastureland and Hayland Interpretations

Cool Season Grasses

Altai wildrye	F
Canada wildrye	G
Crested wheatgrass	G
Green needlegrass	F
Intermediate wheatgrass	F
Meadow bromegrass	F
Newhy hybrid wheatgrass	G
Pubescent wheatgrass	F
Russian wildrye	F
Slender wheatgrass	F
Smooth bromegrass	F
Tall wheatgrass	F
Western wheatgrass	F

Warm Season Grasses

Big bluestem	F
Indiangrass	F
Little bluestem	G
Prairie sandreed	G
Sand bluestem	G
Sideoats grama	F
Switchgrass	F

Legumes

Alfalfa	F
Canada milkvetch	F
Cicer milkvetch	G
Purple prairieclover	F
Sainfoin	F
Sweetclover	F
White prairieclover	G

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

Production Estimates

Production estimates listed here should only be used for making general management recommendations. Onsite production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop**Management Intensity**

	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)
Alfalfa	4000	2000
Alfalfa/Intermediate	3200	1600
Crested wheatgrass	2800	1400
Intermediate wheatgrass	2800	1600
Sand bluestem	3800	2000

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0001

Growth Curve Name: Alfalfa

Growth Curve Description: Alfalfa, MLRAs 107, 102B, 63B, 66, 65

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	30	25	20	15	5	0	0	0

Growth Curve Number: SD0004
Growth Curve Name: Cool season grass
Growth Curve Description: Cool season grass, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Growth Curve Number: SD0003
Growth Curve Name: Irrigated Alfalfa
Growth Curve Description: Irrigated Alfalfa, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	25	20	15	10	0	0	0

Soil Limitations

Soil blowing is a severe hazard during stand establishment or renovation of forage stands on the soils of this group. Bare areas where livestock concentrate are also susceptible. Production potential is low to moderate due to the low available water capacity and droughtiness of these soils. Also, these soils are typically low in native fertility and have reduced capacity to supply plant nutrients. Species choices are somewhat limited for pasture and hayland for these same reasons.

Management Interpretations

The impact on yields of the low available water capacity of these soils can be reduced by selecting forage species that are highly tolerant to periods of drought and inadequate soil moisture and can grow on coarse soils. Incorporate wind erosion control practices during stand establishment. Properly locating facilitating practices such as fences, lanes, and water developments can help control livestock movement, reduce trailing perpendicular to steeper slopes, evenly distribute grazing pressure, and reduce bare areas.

FSG Documentation**Similar FSG's:**

FSG ID	FSG Narrative
G063AY130SD	Very Droughty Loam soils have finer textures than sands.

Inventory Data References

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
 Natural Resources Conservation Service (NRCS) National Water and Climate Center data
 USDA Plant Hardiness Zone Maps,
 National Soil Survey Information System (NASIS) for soil surveys in South Dakota counties in MLRA 63A
 South Dakota NRCS South Dakota Technical Guides
 NRCS National Range and Pasture Handbook
 Various South Dakota Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation

This site has been correlated with the following states: South Dakota

Forage Suitability Group Approval

Original Author: Tim Nordquist
Original Date: 4/5/02
Approval by: Dave Schmidt
Approval Date: 7/17/03